Sherburne County, Minnesota

[Data apply to the entire extent of the map unit within the survey area. Map unit and soil properties for a specific parcel of land may vary somewhat and should be determined by onsite investigation]

### 7A--Hubbard loamy sand, 0 to 2 percent slopes

#### Hubbard

Extent: 95 percent of the unit Soil loss tolerance (T factor): 5

Landform(s): outwash plains, stream terraces Wind erodibility group (WEG): 2

Slope gradient: 0 to 2 percent

Wind erodibility index (WEI): 134

Parent material: outwash

Kw factor (surface layer) .15

Restrictive feature(s): greater than 60 inches

Land capability, nonirrigated: 4s

Flooding: none

Hydric soil: no

Ponding: none

Hydrologic group: A

Drainage class: excessively drained Potential for frost action: low

Representative soil profile:	Texture	Permeability	Available water capacity	рН
Ap,AB 0 to 20 in	loamy sand	rapid	1.61 to 2.41 in	5.1 to 7.3
Bw 20 to 32 in	loamy sand	rapid	0.35 to 0.83 in	5.1 to 7.3
BC,C 32 to 80 in	sand	rapid	1.44 to 3.36 in	5.6 to 7.8

## 7B--Hubbard loamy sand, 2 to 6 percent slopes

#### Hubbard

Extent: 95 percent of the unit Soil loss tolerance (T factor): 5

Landform(s): hills on outwash plains, hills on stream terraces

Wind erodibility group (WEG): 2

Slope gradient: 2 to 6 percent

Wind erodibility index (WEI): 134

Parent material: outwash

Kw factor (surface layer) .15

Restrictive feature(s): greater than 60 inches

Land capability, nonirrigated: 4s

Flooding: none Hydric soil: no
Ponding: none Hydrologic group: A

Drainage class: excessively drained Potential for frost action: low

Representative so	il profile:	Texture	Permeability	Available water capacity	рН
Ap,A 0 to	18 in lo	pamy sand	rapid	1.45 to 2.17 in	5.1 to 7.3
Bw 18 to	23 in lo	pamy sand	rapid	0.14 to 0.33 in	5.1 to 7.3
BC,C 23 to	80 in sa	and	rapid	1.71 to 4.00 in	5.6 to 7.8



Sherburne County, Minnesota

## 7C--Hubbard loamy sand, 6 to 12 percent slopes

#### Hubbard

Extent: 95 percent of the unit Soil loss tolerance (T factor): 5

Landform(s): hills on outwash plains, hills on stream terraces Wind erodibility group (WEG): 2

Slope gradient: 6 to 12 percent Wind erodibility index (WEI): 134

Parent material: outwash Kw factor (surface layer) .15

Restrictive feature(s): greater than 60 inches

Land capability, nonirrigated: 6s

Flooding: none Hydric soil: no
Ponding: none Hydrologic group: A

Drainage class: excessively drained Potential for frost action: low

Representative soil p	rofile:	Texture	Permeability	Available water capacity	рН
Ap,AB 0 to 1	2 in	loamy sand	rapid	0.94 to 1.42 in	5.1 to 7.3
Bw 12 to 3	33 in	coarse sand	rapid	0.64 to 1.49 in	5.1 to 7.3
C 33 to 8	80 in	coarse sand	rapid	1.41 to 3.28 in	5.6 to 7.8

### 32B--Nebish fine sandy loam, 2 to 6 percent slopes

#### Nebish

Extent: 85 percent of the unit

Soil loss tolerance (T factor): 5

Landform(s): hills on moraines

Wind erodibility group (WEG): 3

Slope gradient: 2 to 6 percent Wind erodibility index (WEI): 86
Parent material: till Kw factor (surface layer) .24

Restrictive feature(s): greater than 60 inches

Land capability, nonirrigated: 2e


Flooding: none

Hydric soil: no
Ponding: none

Hydrologic group: B

Drainage class: well drained Potential for frost action: moderate

Representative	soil profile:	Texture	Permeability	Available water capacity	рН
Ap	0 to 5 in	fine sandy loam	moderately rapid	0.67 to 0.92 in	5.6 to 7.3
Bt	5 to 43 in	clay loam	moderate	5.67 to 7.18 in	5.6 to 7.8
C 4	3 to 80 in	loam	moderate	4.07 to 7.03 in	7.4 to 8.4



Sherburne County, Minnesota

## 32C--Nebish fine sandy loam, 6 to 12 percent slopes

#### **Nebish**

Extent: 85 percent of the unit

Landform(s): hills on moraines

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 3

Slope gradient: 6 to 12 percent

Wind erodibility index (WEI): 86

Parent material: till

Kw factor (surface layer) .24

Restrictive feature(s): greater than 60 inches

Land capability, nonirrigated: 3e

Flooding: none

Hydric soil: no

Ponding: none Hydrologic group: B

Drainage class: well drained Potential for frost action: moderate

Representative soil pro	file: Texture	Permeability	capacity	рН
Ap 0 to 7 in	fine sandy loam	moderately rapid	0.92 to 1.28 in	5.6 to 7.3
BE 7 to 11 i	n fine sandy loam	moderately rapid	0.43 to 0.75 in	5.6 to 7.3
Bt 11 to 44 i	n clay loam	moderate	4.96 to 6.28 in	5.6 to 7.8
C 44 to 80 i	n loam	moderate	3.94 to 6.81 in	7.4 to 8.4

### 32D--Nebish fine sandy loam, 12 to 18 percent slopes

#### Nebish

Extent: 85 percent of the unit

Landform(s): hills on moraines

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 3

Slope gradient: 12 to 18 percent

Wind erodibility index (WEI): 86

Parent material: till

Kw factor (surface layer) .24

Restrictive feature(s): greater than 60 inches

Land capability, nonirrigated: 4e


Flooding: none

Hydric soil: no

Hydrologic group: B

Drainage class: well drained Potential for frost action: moderate

Representative	e soil profile:	Texture	Permeability	Available water capacity	рН
Ap	0 to 3 in	fine sandy loam	moderately rapid	0.41 to 0.57 in	5.6 to 7.3
Bt	3 to 22 in	clay loam	moderate	2.83 to 3.59 in	5.6 to 7.8
C	22 to 80 in	loam	moderate	6.37 to 11.00 in	7.4 to 8.4



A collecte contact

Sherburne County, Minnesota

# 32E--Nebish fine sandy loam, 18 to 35 percent slopes

#### **Nebish**

Extent: 85 percent of the unit

Landform(s): hills on moraines

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 3

Slope gradient: 18 to 35 percent

Wind erodibility index (WEI): 86

Parent material: till

Kw factor (surface layer) .24

Restrictive feature(s): greater than 60 inches

Land capability, nonirrigated: 7e

Flooding: none Hydric soil: no
Ponding: none Hydrologic group: B

Drainage class: well drained Potential for frost action: moderate

Representative so	il profile:	Texture	Permeability	Available water capacity	рН
A 0 to	4 in	fine sandy loam	moderately rapid	0.51 to 0.71 in	5.6 to 7.3
E 4 to	13 in	fine sandy loam	moderately rapid	1.00 to 1.72 in	5.6 to 7.3
Bt 13 to	36 in	clay loam	moderate	3.43 to 4.34 in	5.6 to 7.8
C 36 to	80 in	loam	moderate	4.85 to 8.38 in	7.4 to 8.4

Sherburne County, Minnesota

## 38B--Waukon fine sandy loam, 2 to 6 percent slopes

#### Waukon

Extent: 90 percent of the unit

Landform(s): hills on moraines

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 3

Slope gradient: 2 to 6 percent

Wind erodibility index (WEI): 86

Parent material: till

Kw factor (surface layer) .24

Restrictive feature(s): greater than 60 inches

Land capability, nonirrigated: 2e

Flooding: none Hydric soil: no
Ponding: none Hydrologic group: B

Drainage class: well drained Potential for frost action: moderate

Representative soil p	orofile:	Texture	Permeability	capacity	рН
Ap 0 to 8	3 in	fine sandy loam	moderately rapid	1.02 to 1.18 in	6.1 to 7.3
E 8 to 1	12 in	fine sandy loam	moderately rapid	0.43 to 0.75 in	5.6 to 7.3
BE,Bt 12 to 4	43 in	loam	moderate	4.67 to 5.91 in	6.1 to 8.4
Bk 43 to 8	30 in	loam	moderate	5.55 to 7.03 in	7.4 to 8.4

## 75--Bluffton loam, depressional, 0 to 1 percent slopes

#### Bluffton, depressional

Extent: 90 percent of the unit

Landform(s): depressions on moraines

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 5

Slope gradient: 0 to 1 percent

Wind erodibility index (WEI): 56

Parent material: till

Kw factor (surface layer) .28


Restrictive feature(s): greater than 60 inches

Land capability, nonirrigated: 6w

Flooding: none Hydric soil: yes
Ponding: frequent Hydrologic group: B/D

Drainage class: very poorly drained Potential for frost action: high

Representative	soil profile:	Texture	•	Permeability	Available water capacity	рН
A1,A2	0 to 13 in	loam		moderate	2.60 to 3.12 in	5.6 to 7.3
Bg 1	3 to 40 in	loam		moderate	4.07 to 4.62 in	5.6 to 7.3
Cg 4	0 to 80 in	loam		moderately slow	5.96 to 7.56 in	7.4 to 8.4



A. allahla ...atau

Sherburne County, Minnesota

# 125--Beltrami fine sandy loam, 0 to 3 percent slopes

#### **Beltrami**

Extent: 90 percent of the unit

Soil loss tolerance (T factor): 5

Landform(s): moraines

Wind erodibility group (WEG): 3

Slope gradient: 0 to 3 percent

Wind erodibility index (WEI): 86

Parent material: till

Kw factor (surface layer) .24

Restrictive feature(s): greater than 60 inches Land capability, nonirrigated: 1

Flooding: none Hydric soil: no
Ponding: none Hydrologic group: B

Drainage class: somewhat poorly drained Potential for frost action: high

Representative	soil	profile:	Texture	Permeability	Available water capacity	рН
Ap	0 to	6 in	fine sandy loam	moderately rapid	0.77 to 1.06 in	6.1 to 7.3
Bt1	6 to	12 in	loam	moderately rapid	0.65 to 1.12 in	5.6 to 7.3
Bt2,Bt4 1	2 to	48 in	clay loam	moderate	5.43 to 6.88 in	5.6 to 7.8
C 4	8 to	80 in	loam	moderate	4.78 to 6.06 in	7.4 to 8.4



Sherburne County, Minnesota

# 152C--Milaca fine sandy loam, 6 to 12 percent slopes

#### Milaca

Extent: 95 percent of the unit

Landform(s): drumlins

Slope gradient: 6 to 12 percent

Parent material: till

Restrictive feature(s): dense material at 40 to 60 inches

Flooding: none Ponding: none

Drainage class: well drained

Soil loss tolerance (T factor): 4

Wind erodibility group (WEG): 3

Wind erodibility index (WEI): 86

Kw factor (surface layer) .28

Land capability, nonirrigated: 3e

Hydric soil: no

Hydrologic group: C

Potential for frost action: moderate

A collecte contact

Representative soil	profile:	Texture	Permeability	capacity	рН
A 0 to	4 in	fine sandy loam	moderately rapid	0.51 to 0.71 in	5.1 to 6.5
E 4 to	12 in	fine sandy loam	moderately rapid	1.42 to 1.73 in	5.1 to 6.5
Bt 12 to	20 in	fine sandy loam	moderate	0.99 to 1.32 in	5.1 to 6.5
BC 20 to	42 in	sandy loam	slow	1.76 to 2.65 in	5.6 to 7.3
Cd 42 to	80 in	sandy loam	very slow	1.89 to 3.02 in	5.6 to 7.3

Sherburne County, Minnesota

# 152E--Milaca fine sandy loam, 12 to 25 percent slopes

#### Milaca

Extent: 95 percent of the unit

Landform(s): drumlins

Soil loss tolerance (T factor): 4

Wind erodibility group (WEG): 3

Slope gradient: 12 to 25 percent Wind erodibility index (WEI): 86

Parent material: till Kw factor (surface layer) .28

Restrictive feature(s): dense material at 40 to 60 inches Land capability, nonirrigated: 6e

Flooding: none Hydric soil: no
Ponding: none Hydrologic group: C

Drainage class: well drained Potential for frost action: moderate

Representative	soil profile:	Texture	Permeability	Available water capacity	рН
Α	0 to 3 in	fine sandy loam	moderately rapid	0.41 to 0.57 in	5.1 to 6.5
E :	3 to 12 in	fine sandy loam	moderately rapid	1.56 to 1.91 in	5.1 to 6.5
Bt 1:	2 to 20 in	fine sandy loam	moderate	0.99 to 1.32 in	5.1 to 6.5
BC 2	0 to 40 in	sandy loam	slow	1.61 to 2.41 in	5.6 to 7.3
Cd 4	0 to 80 in	sandy loam	very slow	1.99 to 3.18 in	5.6 to 7.3

## 158A--Zimmerman fine sand, 0 to 3 percent slopes

### Zimmerman

Extent: 95 percent of the unit

Landform(s): outwash plains

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 1

Slope gradient: 0 to 3 percent

Wind erodibility index (WEI): 220

Parent material: outwash


Kw factor (surface layer) .17

Restrictive feature(s): greater than 60 inches Land capability, nonirrigated: 4s

Flooding: none Hydric soil: no Hydrologic group: A

Drainage class: excessively drained Potential for frost action: low

Representative	soil profile:	Texture	Permeability	Available water capacity	рН
Ap	0 to 7 in	fine sand	rapid	0.50 to 0.64 in	5.1 to 6.5
E,Bw,E&Bt	7 to 80 in	fine sand	rapid	4.37 to 7.28 in	5.1 to 7.3



This report shows only the major soils in each map unit

Sherburne County, Minnesota

## 158B--Zimmerman fine sand, 3 to 6 percent slopes

#### Zimmerman

Extent: 95 percent of the unit

Landform(s): hills on outwash plains

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 1

Slope gradient: 3 to 6 percent

Wind erodibility index (WEI): 220

Parent material: outwash

Kw factor (surface layer) .17

Restrictive feature(s): greater than 60 inches

Land capability, nonirrigated: 4s

Flooding: none Hydric soil: no
Ponding: none Hydrologic group: A

Drainage class: excessively drained Potential for frost action: low

Representativ	e soil profile:	Texture	Permeability	capacity	рН
Ар	0 to 6 in	fine sand	rapid	0.41 to 0.53 in	5.1 to 6.5
E,Bw,E&Bt	6 to 80 in	fine sand	rapid	4.44 to 7.40 in	5.1 to 7.3

## 158C--Zimmerman fine sand, 6 to 12 percent slopes

#### Zimmerman

Extent: 95 percent of the unit

Landform(s): hills on outwash plains

Wind erodibility group (WEG): 1

Slope gradient: 6 to 12 percent

Wind erodibility index (WEI): 220

Parent material: outwash

Kw factor (surface layer) .17


Restrictive feature(s): greater than 60 inches

Land capability, nonirrigated: 6s

Flooding: noneHydric soil: noPonding: noneHydrologic group: A

Drainage class: excessively drained Potential for frost action: low

Representative	soil profile:	Texture	Permeability	Available water capacity	рН
Ap (	0 to 6 in	fine sand	rapid	0.41 to 0.53 in	5.1 to 6.5
E,Bw,E&Bt	6 to 80 in	fine sand	rapid	4.44 to 7.40 in	5.1 to 7.3



Sherburne County, Minnesota

### 158E--Zimmerman fine sand, 12 to 25 percent slopes

#### Zimmerman

Extent: 95 percent of the unit

Landform(s): hills on outwash plains

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 1

Slope gradient: 12 to 25 percent

Wind erodibility index (WEI): 220

Parent material: outwash

Kw factor (surface layer) .17

Restrictive feature(s): greater than 60 inches

Land capability, nonirrigated: 6s

Flooding: none Hydric soil: no
Ponding: none Hydrologic group: A

Drainage class: excessively drained Potential for frost action: low

Representativ	e soil profile:	Texture	Permeability	capacity	рН
Α	0 to 3 in	fine sand	rapid	0.22 to 0.28 in	5.1 to 6.5
E,Bw,E&Bt	3 to 80 in	fine sand	rapid	4.61 to 7.68 in	5.1 to 7.3

## 161--Isanti fine sandy loam, depressional, 0 to 1 percent slopes

#### Isanti, depressional

Extent: 95 percent of the unit

Landform(s): depressions on outwash plains

Soil loss tolerance (T factor): 3

Wind erodibility group (WEG): 3

Slope gradient: 0 to 1 percent

Wind erodibility index (WEI): 86

Parent material: outwash

Kw factor (surface layer) .20

Restrictive feature(s): greater than 60 inches

Land capability, nonirrigated: 6w

Flooding: none

Hydric soil: yes

Ponding: frequent

Hydrologic group: B/D

Drainage class: very poorly drained Potential for frost action: moderate

Representative	soil profile:	Texture	Permeability	Available water capacity	рН
A1,A3	0 to 16 in	fine sandy loam	rapid	2.10 to 2.91 in	5.1 to 6.5
Bg 1	6 to 28 in	loamy fine sand	rapid	0.71 to 0.94 in	5.1 to 6.5
Cg 2	28 to 80 in	fine sand	rapid	2.60 to 3.64 in	5.6 to 6.5



Sherburne County, Minnesota

# 162--Lino loamy fine sand, 0 to 2 percent slopes

#### Lino

Extent: 95 percent of the unit

Landform(s): outwash plains

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 2

Slope gradient: 0 to 2 percent

Wind erodibility index (WEI): 134

Parent material: outwash

Kw factor (surface layer) .17

Restrictive feature(s): greater than 60 inches

Land capability, nonirrigated: 3s

Flooding: none Hydric soil: no
Ponding: none Hydrologic group: A

Drainage class: somewhat poorly drained Potential for frost action: moderate

Representative :	soil profile:	Texture	Permeability	Available water capacity	рН
Ap 0	) to 8 in	loamy fine sand	rapid	0.79 to 0.94 in	5.1 to 6.0
Bw 8	3 to 38 in	loamy fine sand	rapid	1.80 to 2.39 in	5.1 to 6.0
Ca 38	3 to 80 in	fine sand	rapid	2.11 to 2.95 in	5.1 to 6.5

Sherburne County, Minnesota

## 164A--Mora loam, 0 to 3 percent slopes

#### Mora

Extent: 95 percent of the unit

Landform(s): drumlins

Slope gradient: 0 to 3 percent

Parent material: till

Restrictive feature(s): dense material at 40 to 60 inches

Flooding: none Ponding: none

Drainage class: somewhat poorly drained

Soil loss tolerance (T factor): 4

Wind erodibility group (WEG): 5

Wind erodibility index (WEI): 56

Kw factor (surface layer) .28

w lactor (surface layer) .26

Land capability, nonirrigated: 2s

Hydric soil: no

Hydrologic group: C

Representative	soi	profile:	Texture	Permeability	Available water capacity	рН
Ap	0 to	6 in	loam	moderate	1.00 to 1.18 in	5.1 to 6.5
E	6 to	15 in	sandy loam	moderately rapid	1.27 to 1.72 in	5.1 to 6.5
BE,Bt 1	5 to	36 in	sandy loam	moderate	3.13 to 3.96 in	5.6 to 6.5
BC 3	6 to	48 in	sandy loam	slow	0.98 to 1.46 in	5.6 to 7.3
Cd 4	8 to	80 in	sandy loam	very slow	1.59 to 2.55 in	5.6 to 7.3

Sherburne County, Minnesota

# 165--Parent loam, 0 to 2 percent slopes

#### **Parent**

Extent: 90 percent of the unit

Landform(s): flats on interdrumlins

Slope gradient: 0 to 2 percent

Parent material: till

Restrictive feature(s): dense material at 40 to 60 inches

Flooding: none Ponding: none

Drainage class: poorly drained

Soil loss tolerance (T factor): 4
Wind erodibility group (WEG): 5
Wind erodibility index (WEI): 56
Kw factor (surface layer) .32
Land capability, nonirrigated: 2w

Hydric soil: yes

Hydrologic group: C/D

Representative soil profile:	Texture	Permeability	Available water capacity	рН
A1,A2 0 to 15 in	loam	moderate	2.99 to 3.29 in	5.6 to 7.3
Bg 15 to 33 in	loam	moderate	2.17 to 3.08 in	5.6 to 7.3
BC 33 to 40 in	sandy loam	moderately slow	0.57 to 0.85 in	6.1 to 7.3
Cd 40 to 80 in	sandy loam	very slow	1.99 to 3.18 in	6.1 to 7.3

Sherburne County, Minnesota

# 166--Ronneby loam, 0 to 2 percent slopes

### Ronneby

Extent: 90 percent of the unit Landform(s): interdrumlins Slope gradient: 0 to 2 percent

Parent material: till

Restrictive feature(s): dense material at 40 to 60 inches

Flooding: none Ponding: none

Drainage class: somewhat poorly drained

Soil loss tolerance (T factor): 4
Wind erodibility group (WEG): 5
Wind erodibility index (WEI): 56
Kw factor (surface layer) .32
Land capability, nonirrigated: 2w

Hydric soil: no Hydrologic group: C

Representative	soil profile:	Texture	Permeability	Available water capacity	рН
A	0 to 4 in	loam	moderate	0.71 to 0.91 in	5.1 to 6.5
E	4 to 12 in	fine sandy loam	moderately rapid	0.94 to 1.50 in	5.1 to 6.5
BE,Bt 1	2 to 45 in	fine sandy loam	moderate	3.97 to 6.28 in	5.6 to 6.5
BC 4	15 to 56 in	fine sandy loam	slow	0.88 to 1.32 in	5.6 to 7.3
Cd 5	66 to 80 in	fine sandy loam	very slow	1.20 to 1.92 in	5.6 to 7.3

Sherburne County, Minnesota

## 169B--Braham loamy fine sand, 3 to 6 percent slopes

#### **Braham**

Extent: 90 percent of the unit

Landform(s): hills on moraines

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 2

Slope gradient: 3 to 6 percent

Wind erodibility index (WEI): 134

Parent material: outwash over till

Kw factor (surface layer) .17

Restrictive feature(s): greater than 60 inches

Land capability, nonirrigated: 3s

Flooding: none Hydric soil: no
Ponding: none Hydrologic group: B

Drainage class: moderately well drained Potential for frost action: low

Representative s	oil profile:	Texture	Permeability	capacity	рН
Ap 0	to 9 in	loamy fine sand	rapid	0.91 to 1.09 in	5.6 to 7.3
E 9	to 21 in	loamy fine sand	rapid	0.94 to 1.18 in	5.6 to 7.3
2Bt 21	to 46 in	clay loam	moderate	3.78 to 4.54 in	5.1 to 7.3
2Bk 46	to 80 in	loam	moderate	5.08 to 6.09 in	7.4 to 8.4



A collecte contact

Sherburne County, Minnesota

# 169C--Braham loamy fine sand, 6 to 12 percent slopes

#### **Braham**

Extent: 90 percent of the unit

Landform(s): hills on moraines

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 2

Slope gradient: 6 to 12 percent

Wind erodibility index (WEI): 134

Parent material: outwash over till

Kw factor (surface layer) .17

Restrictive feature(s): greater than 60 inches

Land capability, nonirrigated: 4e

Flooding: none Hydric soil: no
Ponding: none Hydrologic group: B

Drainage class: well drained Potential for frost action: low

Representative soil profile:	Texture	Permeability	Available water capacity	рН
Ap 0 to 8 in	loamy fine sand	rapid	0.79 to 0.94 in	5.6 to 7.3
E 8 to 28 in	loamy sand	rapid	1.61 to 2.01 in	5.6 to 7.3
2Bt 28 to 48 in	clay loam	moderate	3.01 to 3.61 in	5.1 to 7.3
2Bk 48 to 80 in	loam	moderate	4.78 to 5.74 in	7.4 to 8.4

Sherburne County, Minnesota

# 169D--Braham loamy fine sand, 12 to 18 percent slopes

#### **Braham**

Extent: 90 percent of the unit

Landform(s): hills on moraines

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 2

Slope gradient: 12 to 18 percent

Wind erodibility index (WEI): 134

Parent material: outwash over till

Kw factor (surface layer) .17

Restrictive feature(s): greater than 60 inches

Land capability, nonirrigated: 4e

Flooding: none Hydric soil: no
Ponding: none Hydrologic group: B

Drainage class: well drained Potential for frost action: low

Representative so	oil profile:	Texture	Permeability	capacity	pН
Ap 0	to 8 in	loamy fine sand	rapid	0.79 to 0.94 in	5.6 to 7.3
E 8	to 32 in	loamy sand	rapid	1.92 to 2.40 in	5.6 to 7.3
2Bt 32	to 55 in	clay loam	moderate	3.48 to 4.18 in	5.1 to 7.3
2Bk 55	to 80 in	loam	moderate	3.72 to 4.46 in	7.4 to 8.4



Sherburne County, Minnesota

# 204B--Cushing fine sandy loam, 2 to 8 percent slopes

## Cushing

Extent: 90 percent of the unit

Landform(s): hills on moraines

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 3

Slope gradient: 2 to 8 percent

Wind erodibility index (WEI): 86

Parent material: till

Kw factor (surface layer) .24

Restrictive feature(s): greater than 60 inches

Land capability, nonirrigated: 2e

Flooding: none

Hydric soil: no
Ponding: none

Hydrologic group: B

Drainage class: well drained Potential for frost action: moderate

Representative so	oil profile:	Texture	Permeability	Available water capacity	рН
A 0 1	o 6 in	fine sandy loam	moderate	0.59 to 1.30 in	5.1 to 7.8
E,B/E 6	o 22 in	fine sandy loam	moderate	1.61 to 3.55 in	5.1 to 7.8
Bt,BC 22	o 44 in	clay loam	moderate	2.20 to 4.19 in	5.1 to 7.8
C 44	o 80 in	loam	moderately slow	3.22 to 6.81 in	5.1 to 8.4



Sherburne County, Minnesota

## 204C--Cushing fine sandy loam, 8 to 15 percent slopes

### Cushing

Extent: 95 percent of the unit

Landform(s): hills on moraines

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 3

Slope gradient: 8 to 15 percent

Wind erodibility index (WEI): 86

Parent material: till

Kw factor (surface layer) .24

Restrictive feature(s): greater than 60 inches

Land capability, nonirrigated: 3e

Flooding: none Hydric soil: no
Ponding: none Hydrologic group: B

Drainage class: well drained Potential for frost action: moderate

Representative soil profile:	Texture	Permeability	capacity	рН
Ap 0 to 7 in	fine sandy loam	moderate	0.71 to 1.56 in	5.1 to 7.8
E 7 to 21 in	fine sandy loam	moderate	1.38 to 3.03 in	5.1 to 7.8
Bt 21 to 44 in	clay loam	moderate	2.32 to 4.41 in	5.1 to 7.8
C 44 to 80 in	sandy loam	moderately slow	3.22 to 6.81 in	5.1 to 8.4

### 258B--Sandberg loamy coarse sand, 1 to 6 percent slopes

#### Sandberg

Extent: 95 percent of the unit

Landform(s): hills on stream terraces

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 2

Slope gradient: 1 to 6 percent

Wind erodibility index (WEI): 134

Parent material: outwash

Kw factor (surface layer) .15

Restrictive feature(s): greater than 60 inches

Land capability, nonirrigated: 4s

Flooding: none

Hydric soil: no

Ponding: none

Hydrologic group: A

Drainage class: excessively drained Potential for frost action: low

profile:	Texture	Permeability	capacity	рН
14 in	loamy coarse sand	rapid	1.42 to 1.70 in	5.6 to 7.8
32 in	gravelly coarse sand	rapid	0.53 to 1.77 in	6.1 to 7.8
80 in	sand	very rapid	0.96 to 2.88 in	7.4 to 8.4
	14 in 32 in	14 in loamy coarse sand 32 in gravelly coarse sand	14 in loamy coarse sand rapid 32 in gravelly coarse sand rapid	profile: Texture Permeability capacity  14 in loamy coarse sand rapid 1.42 to 1.70 in gravelly coarse sand rapid 0.53 to 1.77 in



A collecte contact

Available water

Sherburne County, Minnesota

## 258C--Sandberg loamy coarse sand, 6 to 12 percent slopes

### Sandberg

Extent: 95 percent of the unit

Landform(s): hills on stream terraces

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 2

Slope gradient: 6 to 12 percent

Wind erodibility index (WEI): 134

Parent material: outwash

Kw factor (surface layer) .15

Restrictive feature(s): greater than 60 inches

Land capability, nonirrigated: 6s

Flooding: none Hydric soil: no
Ponding: none Hydrologic group: A

Drainage class: excessively drained Potential for frost action: low

Representative soil	profile:	Texture	Permeability	Available water capacity	pН
Ap 0 to	11 in	loamy coarse sand	rapid	1.10 to 1.32 in	5.6 to 7.8
Bw 11 to	26 in	coarse sand	rapid	0.45 to 1.50 in	6.1 to 7.8
C 26 to	80 in	coarse sand	very rapid	1.08 to 3.24 in	7.4 to 8.4

## 258E--Sandberg loamy coarse sand, 12 to 35 percent slopes

#### Sandberg

Extent: 95 percent of the unit Soil loss tolerance (T factor): 5

Landform(s): escarpments on stream terraces, hills on stream Wind erodibility group (WEG): 2

terraces

Slope gradient: 12 to 35 percent

Wind erodibility index (WEI): 134

Parent material: outwash

Kw factor (surface layer) .15


Restrictive feature(s): greater than 60 inches

Land capability, nonirrigated: 7e

Flooding: none Hydric soil: no Ponding: none Hydrologic group: A

Drainage class: excessively drained Potential for frost action: low

Representative se	oil profile:	Texture	Permeability	capacity	рН
A 0	to 11 in	loamy coarse sand	rapid	1.10 to 1.32 in	5.6 to 7.8
Bw 11	to 27 in	coarse sand	rapid	0.48 to 1.61 in	6.1 to 7.8
C 27	to 80 in	gravelly coarse sand	very rapid	1.06 to 3.17 in	7.4 to 8.4



This report shows only the major soils in each map unit

L Available water

Sherburne County, Minnesota

### 260--Duelm loamy sand, 0 to 2 percent slopes

#### **Duelm**

Extent: 95 percent of the unit

Soil loss tolerance (T factor): 5

Landform(s): outwash plains, stream terraces

Wind erodibility group (WEG): 2

Slope gradient: 0 to 2 percent

Wind erodibility index (WEI): 134

Parent material: outwash

Kw factor (surface layer) .17

Restrictive feature(s): greater than 60 inches

Land capability, nonirrigated: 4s

Flooding: none

Hydric soil: no
Ponding: none

Hydrologic group: A

Drainage class: moderately well drained Potential for frost action: moderate

Representative soil pr	rofile:	Texture	Permeability	Available water capacity	pН
Ap,AB 0 to 16	6 in loamy sand		rapid	1.29 to 1.94 in	5.6 to 7.3
Bw 16 to 30	0 in coarse sand		rapid	0.83 to 1.52 in	5.1 to 7.3
C 30 to 80	0 in coarse sand		rapid	1.00 to 3.50 in	5.6 to 7.8

### 261--Isan sandy loam, depressional, 0 to 1 percent slopes

#### Isan, depressional

Extent: 95 percent of the unit

Soil loss tolerance (T factor): 3

Landform(s): depressions on outwash plains, depressions on

Wind erodibility group (WEG): 3

stream terraces

Slope gradient: 0 to 1 percent

Wind erodibility index (WEI): 86

Parent material: outwash

Kw factor (surface layer) .20

Restrictive feature(s): greater than 60 inches


Land capability, nonirrigated: 6w

Flooding: none Hydric soil: yes

Ponding: frequent Hydrologic group: B/D

Drainage class: very poorly drained Potential for frost action: moderate

Representative soil	profile:	Texture	Permeability	capacity	pН
A 0 to	14 in	sandy loam	moderately rapid	1.42 to 2.13 in	5.6 to 7.3
AB,Bg 14 to	34 in	loamy sand	rapid	1.18 to 1.97 in	5.1 to 6.5
Cg 34 to	80 in	coarse sand	rapid	1.84 to 2.76 in	5.6 to 7.3



L Available water

Sherburne County, Minnesota

## 325--Prebish fine sandy loam, depressional, 0 to 1 percent slopes

### Prebish, depressional

Extent: 95 percent of the unit

Landform(s): depressions on interdrumlins

Soil loss tolerance (T factor): 4

Wind erodibility group (WEG): 3

Slope gradient: 0 to 1 percent Wind erodibility index (WEI): 86

Parent material: till Kw factor (surface layer) .20
Restrictive feature(s): dense material at 40 to 60 inches Land capability, nonirrigated: 6w

Flooding: none Hydric soil: yes
Ponding: frequent Hydrologic group: C/D

Drainage class: very poorly drained Potential for frost action: high

Representative soil	profile:	Texture	Permeability	Available water capacity	рН
A1,A2 0 to	15 in	fine sandy loam	moderately rapid	2.39 to 2.69 in	5.6 to 7.3
Bg,BC 15 to	41 in	sandy loam	moderate	3.64 to 4.16 in	5.6 to 7.3
Cd 41 to	80 in	fine sandy loam	very slow	1.95 to 3.12 in	5.6 to 7.3

### 341--Arvilla sandy loam, 0 to 2 percent slopes

#### Arvilla

Extent: 95 percent of the unit

Landform(s): stream terraces

Slope gradient: 0 to 2 percent

Parent material: outwash

Restrictive feature(s): greater than 60 inches

Soil loss tolerance (T factor): 3

Wind erodibility group (WEG): 3

Wind erodibility index (WEI): 86

Kw factor (surface layer) .20

Land capability, nonirrigated: 3s

Flooding: none Hydric soil: no
Ponding: none Hydrologic group: B

Drainage class: somewhat excessively drained Potential for frost action: low

Representativ	e soil profile:	Texture	Permeability	Available water capacity	рН
Ap,A	0 to 14 in	sandy loam	moderately rapid	1.84 to 2.13 in	6.1 to 7.3
Bw	14 to 20 in	sandy loam	moderately rapid	0.65 to 0.83 in	6.1 to 7.3
2Bw,2Bk,2C	20 to 80 in	gravelly coarse sand	rapid	1.20 to 2.99 in	7.4 to 8.4



Sherburne County, Minnesota

## 346--Talmoon loam, 0 to 2 percent slopes

#### **Talmoon**

Extent:90 percent of the unitSoil loss tolerance (T factor):5Landform(s):swales on morainesWind erodibility group (WEG):5Slope gradient:0 to 2 percentWind erodibility index (WEI):56Parent material:tillKw factor (surface layer).28

Restrictive feature(s): greater than 60 inches Land capability, nonirrigated: 2w

Flooding: none Hydric soil: yes
Ponding: none Hydrologic group: B/D

Drainage class: poorly drained Potential for frost action: high

Representative soil	profile:	Texture	Permeability	capacity	рН
Ap 0 to	7 in	loam	moderate	1.42 to 1.56 in	5.1 to 7.3
Eg 7 to	12 in	fine sandy loam	moderate	0.61 to 1.04 in	5.1 to 7.3
Btg 12 to	32 in	clay loam	moderately slow	3.21 to 3.81 in	5.6 to 7.3
Bk 32 to	80 in	loam	moderately slow	7.20 to 9.13 in	7.4 to 8.4

### 373--Renshaw loam, 0 to 3 percent slopes

#### Renshaw

Extent: 95 percent of the unit

Landform(s): stream terraces

Solope gradient: 0 to 3 percent

Parent material: outwash

Restrictive feature(s): greater than 60 inches

Solope Solope Wind erodibility group (WEG): 6

Wind erodibility index (WEI): 48

Kw factor (surface layer) .28

Land capability, nonirrigated: 3s

Flooding: none Hydric soil: no
Ponding: none Hydrologic group: B

Drainage class: somewhat excessively drained Potential for frost action: low

Representativ	ve soil profile:	Texture	Permeability	Available water capacity	рН
Ар	0 to 9 in	loam	moderate	1.63 to 1.81 in	6.0 to 7.3
Bw	9 to 15 in	loam	moderately rapid	0.65 to 1.06 in	6.0 to 7.3
2Bw,2C	15 to 80 in	gravelly coarse sand	very rapid	1.95 to 3.90 in	6.6 to 8.4



This report shows only the major soils in each map unit

A collecte contact

Sherburne County, Minnesota

### 454B--Mahtomedi loamy coarse sand, 1 to 6 percent slopes

#### Mahtomedi

Extent: 95 percent of the unit

Landform(s): hills on moraines

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 2

Slope gradient: 1 to 6 percent

Wind erodibility index (WEI): 134

Parent material: outwash

Kw factor (surface layer) .15

Restrictive feature(s): greater than 60 inches

Land capability, nonirrigated: 4s

Flooding: none Hydric soil: no
Ponding: none Hydrologic group: A

Drainage class: excessively drained Potential for frost action: low

Representative	soil profile:	Texture	Permeability	Available water capacity	рН
Ap	0 to 9 in	loamy coarse sand	rapid	0.91 to 1.09 in	5.1 to 6.5
Bw	9 to 36 in	gravelly coarse sand	rapid	1.34 to 1.87 in	5.1 to 6.5
BC,C 3	6 to 80 in	coarse sand	rapid	1.76 to 3.97 in	5.1 to 7.3

## 454C--Mahtomedi loamy coarse sand, 6 to 15 percent slopes

#### Mahtomedi

Extent: 95 percent of the unit

Landform(s): hills on moraines

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 2

Slope gradient: 6 to 15 percent

Wind erodibility index (WEI): 134

Parent material: outwash

Kw factor (surface layer) .15

Restrictive feature(s): greater than 60 inches

Land capability, nonirrigated: 6s

Hydric soil: no

Ponding: none Hydrologic group: A

Drainage class: excessively drained Potential for frost action: low

Representative soil profile:		Texture	Permeability	Available water capacity	рН
Ap	0 to 3 in	loamy coarse sand	rapid	0.31 to 0.38 in	5.1 to 6.5
Bw	3 to 17 in	gravelly sand	rapid	0.69 to 0.96 in	5.1 to 6.5
C	17 to 80 in	sand	rapid	2.52 to 5.67 in	5.1 to 7.3



Sherburne County, Minnesota

## 540--Seelyeville muck, 0 to 1 percent slopes

### Seelyeville

Extent: 95 percent of the unit Soil loss tolerance (T factor): 3

Landform(s): depressions on moraines, depressions on outwash plains, depressions on stream terraces

Slope gradient: 0 to 1 percent

Wind erodibility index (WEI): 134

Parent material: organic material

Kw factor (surface layer) .02

Restrictive feature(s): greater than 60 inches

Land capability, nonirrigated: 6w

Flooding: none Hydric soil: yes
Ponding: frequent Hydrologic group: A/D

Drainage class: very poorly drained Potential for frost action: high

Representative soil profile:

Oa1 -- 0 to 10 in muck

Oa2,Oa5 -- 10 to 80 in muck

Texture

Permeability

Available water capacity

pH

moderately rapid 3.44 to 4.43 in moderately rapid 24.53 to 31.54 in

# 543--Markey muck, 0 to 1 percent slopes

#### Markey

Extent: 90 percent of the unit

Soil loss tolerance (T factor): 2

Landform(s): depressions on outwash plains, depressions on Wind erodibility group (WEG): 2 stream terraces

Slope gradient: 0 to 1 percent Wind erodibility index (WEI): 134

Parent material: organic material over outwash

Kw factor (surface layer) .02

Restrictive feature(s): greater than 60 inches

Land capability, nonirrigated: 6w

Flooding: none Hydric soil: yes
Ponding: frequent Hydrologic group: A/D

Drainage class: very poorly drained Potential for frost action: high

Representative soil profile:	Texture	Permeability	capacity	рН
Oa 0 to 36 in	muck	moderately rapid	12.54 to 16.12 in	
2A 36 to 42 in	loamy sand	rapid	0.19 to 0.50 in	
2Cg 42 to 80 in	sand	rapid	1.13 to 3.02 in	



Available water

Sherburne County, Minnesota

## 544--Cathro muck, 0 to 1 percent slopes

#### Cathro

Extent: 95 percent of the unit Soil loss tolerance (T factor): 2

Landform(s): depressions on moraines Wind erodibility group (WEG): 2

Slope gradient: 0 to 1 percent

Wind erodibility index (WEI): 134

Parent material: organic material over till

Kw factor (surface layer) .02

Restrictive feature(s): greater than 60 inches

Land capability, nonirrigated: 6w

Flooding: none Hydric soil: yes
Ponding: frequent Hydrologic group: A/D

Drainage class: very poorly drained Potential for frost action: high

Representative soil profile:	Texture	Permeability	Available water capacity	рН
Oa1,Oa2 0 to 30 in	muck	moderately rapid	13.46 to 16.46 in	
2A 30 to 38 in	loam	moderate	0.87 to 1.73 in	
2Cq 38 to 80 in	loam	moderate	4.63 to 9.27 in	

### 565--Eckvoll loamy fine sand, 0 to 3 percent slopes

#### **Eckvoll**

Extent: 90 percent of the unit

Landform(s): moraines

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 2

Slope gradient: 0 to 3 percent

Wind erodibility index (WEI): 134

Parent material: outwash over till

Restrictive feature(s): greater than 60 inches

Land capability, nonirrigated: 3s

Flooding: none Hydric soil: no
Ponding: none Hydrologic group: B

Drainage class: moderately well drained Potential for frost action: high

Representative soil profile:	Texture	Permeability	Available water capacity	рН
Ap 0 to 9 in	loamy fine sand	rapid	0.91 to 1.09 in	6.1 to 7.3
E 9 to 24 in	fine sand	rapid	0.90 to 1.20 in	6.1 to 7.3
2Bt 24 to 45 in	loam	moderate	3.34 to 3.76 in	6.1 to 7.3
2C 45 to 80 in	loam	moderate	5.96 to 6.66 in	7.4 to 8.4



This report shows only the major soils in each map unit

Sherburne County, Minnesota

## 567--Verndale sandy loam, 0 to 2 percent slopes

#### Verndale

Extent: 95 percent of the unit Soil loss tolerance (T factor): 3

Landform(s): outwash plains, stream terraces Wind erodibility group (WEG): 3

Slope gradient: 0 to 2 percent

Wind erodibility index (WEI): 86

Parent material: outwash

Restrictive feature(s): greater than 60 inches

Wind erodibility index (WEI): 86

Kw factor (surface layer) .20

Land capability, nonirrigated: 3s

Flooding: none Hydric soil: no
Ponding: none Hydrologic group: B

Drainage class: somewhat excessively drained Potential for frost action: low

Representative soil profile:	Texture	Permeability	capacity	рН
Ap 0 to 10 in	sandy loam	moderately rapid	1.28 to 1.67 in	5.1 to 7.3
Bt 10 to 19 in	sandy loam	moderate	1.27 to 1.63 in	5.1 to 7.3
2Bw 19 to 28 in	sand	rapid	0.54 to 0.72 in	5.1 to 7.3
2C 28 to 80 in	sand	rapid	1.04 to 3.12 in	5.1 to 7.3

### 623A--Pierz sandy loam, 0 to 2 percent slopes

### **Pierz**

Extent: 95 percent of the unit

Landform(s): outwash plains

Solope gradient: 0 to 2 percent

Parent material: outwash

Restrictive feature(s): greater than 60 inches

Solope Solope Wind erodibility group (WEG): 3

Wind erodibility index (WEI): 86

Kw factor (surface layer) .20

Land capability, nonirrigated: 3s

Flooding: none


Hydric soil: no

Ponding: none

Hydrologic group: B

Drainage class: well drained Potential for frost action: low

Representati	ve soil profile:	Texture	Permeability	capacity	рН
Ap,A,AB	0 to 16 in	sandy loam	moderately rapid	2.10 to 2.74 in	5.1 to 6.5
Bt	16 to 29 in	sandy loam	moderately rapid	2.08 to 2.60 in	5.1 to 6.5
2C	29 to 80 in	gravelly sand	very rapid	1.02 to 2.03 in	5.1 to 6.5



Available water

Sherburne County, Minnesota

### 623B--Pierz sandy loam, 2 to 6 percent slopes

#### **Pierz**

Extent: 95 percent of the unit

Landform(s): hills on outwash plains

Soil loss tolerance (T factor): 4

Wind erodibility group (WEG): 3

Slope gradient: 2 to 6 percent

Wind erodibility index (WEI): 86

Parent material: outwash

Kw factor (surface layer) .20

Restrictive feature(s): greater than 60 inches

Land capability, nonirrigated: 3s

Flooding: none Hydric soil: no
Ponding: none Hydrologic group: B

Drainage class: well drained Potential for frost action: low

Representative soil profile:		Texture	Permeability	Available water capacity	рН
Ap	0 to 9 in	sandy loam	moderately rapid	1.18 to 1.54 in	5.1 to 6.5
Bt	9 to 22 in	sandy loam	moderately rapid	2.08 to 2.60 in	5.1 to 6.5
2C 2	2 to 80 in	very gravelly coarse sand	very rapid	1.16 to 2.31 in	5.1 to 6.5

### 708--Rushlake coarse sand, 1 to 4 percent slopes

#### Rushlake

Extent: 85 percent of the unit

Landform(s): beaches

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 1

Slope gradient: 1 to 4 percent

Wind erodibility index (WEI): 160

Parent material: outwash

Kw factor (surface layer) .15

Restrictive feature(s): greater than 60 inches Land capability, nonirrigated: 4s

Flooding: none

Hydric soil: no

Hydrologic group: A

Drainage class: moderately well drained Potential for frost action: moderate

Representative soil prof	ile: Texture	Permeability	Available water capacity	рН
A 0 to 9 in	coarse sand	rapid	0.36 to 0.63 in	6.1 to 7.8
C 9 to 80 in	coarse sand	rapid	1.42 to 7.09 in	6.5 to 8.4



Sherburne County, Minnesota

### 730A--Sanburn fine sandy loam, 0 to 2 percent slopes

#### Sanburn

Extent: 90 percent of the unit

Landform(s): outwash plains, moraines

Soil loss tolerance (T factor): 3

Wind erodibility group (WEG): 3

Slope gradient: 0 to 2 percent

Wind erodibility index (WEI): 86

Parent material: outwash

Restrictive feature(s): greater than 60 inches

Land capability, nonirrigated: 3s

Flooding: none Hydric soil: no
Ponding: none Hydrologic group: B

Drainage class: somewhat excessively drained Potential for frost action: low

Representative soil profile:		e soil profile:	Texture	Permeability	capacity	pН
	Ар	0 to 6 in	fine sandy loam	moderately rapid	0.71 to 0.89 in	5.1 to 6.5
	Bt	6 to 18 in	sandy loam	moderately rapid	0.85 to 1.46 in	5.1 to 6.5
	2Bt,2BC,2C	18 to 80 in	gravelly coarse sand	rapid	1.24 to 2.47 in	5.1 to 6.5

## 730B--Sanburn fine sandy loam, 2 to 6 percent slopes

#### Sanburn

Extent: 90 percent of the unit

Landform(s): hills on outwash plains, hills on moraines

Slope gradient: 2 to 6 percent

Parent material: outwash

Restrictive feature(s): greater than 60 inches

Flooding: none

Soil loss tolerance (T factor): 3

Wind erodibility group (WEG): 3

Wind erodibility index (WEI): 86

Kw factor (surface layer) .20


Land capability, nonirrigated: 3e

Hydric soil: no

Ponding: none Hydrologic group: B

Drainage class: somewhat excessively drained Potential for frost action: low

Representative soil profile:		Texture	Permeability	Available water capacity	рН
Ap	0 to 5 in	fine sandy loam	moderately rapid	0.61 to 0.77 in	5.1 to 6.5
Bt	5 to 19 in	sandy loam	moderately rapid	0.96 to 1.65 in	5.1 to 6.5
2BC,2C 1	9 to 80 in	gravelly coarse sand	rapid	1.22 to 2.44 in	5.1 to 6.5



Sherburne County, Minnesota

# 732B--Bushville fine sand, 2 to 6 percent slopes

#### **Bushville**

Extent: 95 percent of the unit

Landform(s): drumlins

Slope gradient: 2 to 6 percent Parent material: outwash over till

Restrictive feature(s): dense material at 40 to 60 inches

Flooding: none Ponding: none

Drainage class: somewhat poorly drained

Soil loss tolerance (T factor): 4

Wind erodibility group (WEG): 1

Wind erodibility index (WEI): 220

Kw factor (surface layer) .15

A collecte contact

Land capability, nonirrigated: 3s

Hydric soil: no

Hydrologic group: C

Potential for frost action: low

Representative soil profile:		Texture	Permeability	capacity	рН
Ap	0 to 8 in	fine sand	rapid	0.55 to 0.79 in	5.1 to 6.5
E1,E2,E3	8 to 28 in	fine sand	rapid	1.20 to 1.81 in	5.1 to 6.5
2Bt 2	28 to 33 in	fine sandy loam	moderate	0.61 to 0.82 in	5.1 to 6.5
2BC 3	33 to 43 in	fine sandy loam	slow	0.79 to 1.18 in	5.1 to 7.3
2Cd 4	43 to 80 in	sandy loam	very slow	1.85 to 2.96 in	5.6 to 7.3

Sherburne County, Minnesota

# 768--Mosford sandy loam, 0 to 2 percent slopes

#### **Mosford**

Ponding: none

Extent: 95 percent of the unit

Soil loss tolerance (T factor): 3

Landform(s): outwash plains, stream terraces

Wind erodibility group (WEG): 3

Slope gradient: 0 to 2 percent

Wind erodibility index (WEI): 86

Parent material: outwash

Kw factor (surface layer) .20

Restrictive feature(s): greater than 60 inches

Land capability, nonirrigated: 3s

Flooding: none

Hydric soil: no

Drainage class: somewhat excessively drained Potential for frost action: low

Hydrologic group: B

Representative soil profile	Texture	Permeability	Available water capacity	рН
Ap,A 0 to 12 in	sandy loam	moderately rapid	1.54 to 2.13 in	5.1 to 7.3
Bw 12 to 16 in	coarse sandy loam	moderately rapid	0.52 to 0.74 in	5.1 to 7.3
2Bw 16 to 21 in	coarse sand	rapid	0.14 to 0.52 in	5.1 to 7.3
2C 21 to 80 in	sand	rapid	1.18 to 4.13 in	5.1 to 7.8



Sherburne County, Minnesota

# 771--Elkriver fine sandy loam, 0 to 2 percent slopes, rarely flooded

## Elkriver, rarely flooded

Extent: 95 percent of the unit

Landform(s): flood plains

Soll loss tolerance (T factor): 4

Wind erodibility group (WEG): 3

Slope gradient: 0 to 2 percent

Wind erodibility index (WEI): 86

Parent material: alluvium

Kw factor (surface layer) .20

Restrictive feature(s): greater than 60 inches

Land capability, nonirrigated: 2s

Flooding: rare Hydric soil: no
Ponding: none Hydrologic group: B

Drainage class: moderately well drained Potential for frost action: moderate

Representative	soil profile:	Texture	Permeability	capacity	рН
Ap	0 to 10 in	fine sandy loam	moderately rapid	1.57 to 1.97 in	5.1 to 7.3
A1,A3 1	0 to 35 in	fine sandy loam	moderately rapid	3.78 to 5.04 in	5.1 to 7.3
Bw 3	5 to 39 in	fine sandy loam	moderately rapid	0.59 to 0.75 in	5.6 to 7.8
2C 3	9 to 80 in	sand	rapid	0.82 to 4.09 in	5.6 to 7.8



Sherburne County, Minnesota

## 799--Seelyeville and Bowstring soils, 0 to 1 percent slopes, frequently flooded

# Seelyeville, frequently flooded

Extent: 45 percent of the unit

Landform(s): flood plains

Slope gradient: 0 to 1 percent

Parent material: organic material

Restrictive feature(s): greater than 60 inches

Flooding: frequent Ponding: none

Drainage class: very poorly drained

Soil loss tolerance (T factor): 3
Wind erodibility group (WEG): 2
Wind erodibility index (WEI): 134
Kw factor (surface layer) .02
Land capability, nonirrigated: 6w

Hydric soil: yes

Hydrologic group: A/D

Potential for frost action: high

Representative soil profile:	Texture	Permeability	capacity	pН
Oa1 0 to 12 in	muck	moderately rapid	4.13 to 5.31 in	
Oa2,Oa3 12 to 80 in	muck	moderately rapid	23.84 to 30.65 in	

### Bowstring, frequently flooded

Extent: 45 percent of the unit Landform(s): flood plains Slope gradient: 0 to 1 percent

Parent material: organic material and alluvium Restrictive feature(s): greater than 60 inches

Flooding: frequent Ponding: none


Drainage class: very poorly drained

Soil loss tolerance (T factor): 3
Wind erodibility group (WEG): 2

Wind erodibility index (WEI): 134 Kw factor (surface layer) .02 Land capability, nonirrigated: 6w

Hydric soil: yes Hydrologic group: A/D

Representative soil profile:		Texture	Permeability	Available water capacity	рН
Oa1,Oa2	0 to 38 in	muck	moderately rapid	13.23 to 17.01 in	
Cg 3	38 to 47 in	stratified fine sand to fine sandy loam	rapid	0.72 to 1.27 in	
O'a1 4	47 to 80 in	muck	moderately rapid	11.57 to 14.88 in	



Sherburne County, Minnesota

## 1013--Pits, quarry

### Pits, quarry

Extent: 100 percent of the unit Soil loss tolerance (T factor):

Landform(s): stream terraces Wind erodibility group (WEG):

Slope gradient: Wind erodibility index (WEI):

Parent material: bedrock, granite Kw factor (surface layer)

Restrictive feature(s): lithic bedrock at 0 to 4 inches

Land capability, nonirrigated:

Flooding: Hydric soil:
Ponding: Hydrologic group:

Drainage class: Potential for frost action:

Representative soil profile:

Texture

Permeability

Available water capacity pH

# 1015--Udipsamments, cut and fill land

### Udipsamments, cut and fill land

Extent: 100 percent of the unit Soil loss tolerance (T factor): 5

Landform(s): outwash plains, stream terraces, lake plains Wind erodibility group (WEG): 2

Slope gradient: 0 to 6 percent Wind erodibility index (WEI): 134

Parent material: variable sandy material

Kw factor (surface layer)

Restrictive feature(s): greater than 60 inches

Land capability, nonirrigated:

Flooding: none

Hydric soil:

Ponding: none Hydrologic group: A
Drainage class: excessively drained Potential for frost action:

Representative soil profile:

Texture

Permeability

Available water capacity

pH



Sherburne County, Minnesota

# 1016--Udorthents, loamy, cut and fill land

## Udorthents, loamy, cut and fill land

Extent: 100 percent of the unit

Landform(s): moraines

Slope gradient: 0 to 12 percent

Parent material: variable loamy material

Restrictive feature(s): greater than 60 inches

Flooding: none Ponding: none

Drainage class: well drained

Representative soil profile:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated:

Hydric soil:

Hydrologic group: B

Potential for frost action:

Texture Permeability

Available water capacity

рΗ



Sherburne County, Minnesota

## 1028--Udorthents-Pits, gravel, complex

#### **Udorthents**

Extent: 55 percent of the unit Soil loss tolerance (T factor):

Landform(s): moraines, outwash plains, stream terraces Wind erodibility group (WEG):

Slope gradient: 0 to 25 percent

Wind erodibility index (WEI):

Parent material: outwash

Kw factor (surface laver)

Parent material: outwash

Kw factor (surface layer)

Restrictive feature(s): greater than 60 inches

Land capability, nonirrigated:

Hydric soil:

Ponding: Hydrologic group: A

Drainage class: excessively drained Potential for frost action:

Representative soil profile:

Texture

Permeability

Available water capacity

pH

### Pits, gravel

Extent: 45 percent of the unit Soil loss tolerance (T factor):

Landform(s): moraines, outwash plains, stream terraces Wind erodibility group (WEG):

Slope gradient: 0 to 50 percent Wind erodibility index (WEI):

Parent material: sandy and gravelly outwash

Restrictive feature(s): greater than 60 inches

Kw factor (surface layer)

Land capability, nonirrigated:

Flooding: Hydric soil:
Ponding: Hydrologic group:

Drainage class: Potential for frost action:

Representative soil profile:

Texture

Permeability

Available water capacity

pH



Sherburne County, Minnesota

## 1109--Isanti loamy fine sand, 0 to 2 percent slopes

#### Isanti

Extent: 90 percent of the unit

Landform(s): swales on outwash plains

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 2

Slope gradient: 0 to 2 percent

Wind erodibility index (WEI): 134

Parent material: outwash

Kw factor (surface layer) .17

Restrictive feature(s): greater than 60 inches

Land capability, nonirrigated: 3w

Flooding: none Hydric soil: yes
Ponding: none Hydrologic group: A/D

Drainage class: poorly drained Potential for frost action: moderate

Representative soil profile	Texture	Permeability	Available water capacity	рН
Ap,AB 0 to 16 in	loamy fine sand	rapid	1.61 to 1.94 in	5.1 to 6.5
Bg 16 to 23 in	loamy fine sand	rapid	0.40 to 0.54 in	5.1 to 6.5
BCq,Cq 23 to 80 in	fine sand	rapid	2.85 to 4.00 in	5.6 to 6.5

## 1110--Isan sandy loam, 0 to 2 percent slopes

#### Isan

Extent: 90 percent of the unit

Soil loss tolerance (T factor): 3

Landform(s): swales on stream terraces, swales on outwash Wind erodibility group (WEG): 3 plains

Slope gradient: 0 to 2 percent Wind erodibility index (WEI): 86
Parent material: outwash Kw factor (surface layer) .20

Restrictive feature(s): greater than 60 inches

Land capability, nonirrigated: 3w

Flooding: none Hydric soil: yes
Ponding: none Hydrologic group: B/D

Drainage class: poorly drained Potential for frost action: moderate

Representative soil p	orofile:	Texture	Permeability	Available water capacity	рН
Ap,A 0 to	18 in sandy loam		moderately rapid	1.81 to 2.72 in	5.6 to 7.3
AB,Bg 18 to 2	29 in loamy sand		rapid	0.66 to 1.10 in	5.1 to 6.5
Cg 29 to 8	80 in coarse sand		rapid	2.03 to 3.05 in	5.6 to 7.3



Sherburne County, Minnesota

# 1223--Sandberg-Arvilla complex, 0 to 3 percent slopes

#### Sandberg

Extent: 60 percent of the unit

Landform(s): stream terraces

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 2

Slope gradient: 1 to 3 percent

Wind erodibility index (WEI): 134

Parent material: outwash

Kw factor (surface layer) .15

Restrictive feature(s): greater than 60 inches

Land capability, nonirrigated: 4s

Flooding: none Hydric soil: no
Ponding: none Hydrologic group: A

Drainage class: excessively drained Potential for frost action: low

Representative soil profi	le: Texture	Permeability	capacity	рН
Ap 0 to 11 in	loamy coarse sand	rapid	1.10 to 1.32 in	5.6 to 7.8
Bw,BC 11 to 35 in	gravelly coarse sand	rapid	0.72 to 2.40 in	6.1 to 7.8
C 35 to 80 in	gravelly coarse sand	very rapid	0.90 to 2.69 in	7.4 to 8.4

#### Arvilla

Extent: 30 percent of the unit

Landform(s): stream terraces

Slope gradient: 0 to 2 percent

Parent material: outwash

Restrictive feature(s): greater than 60 inches

Soil loss tolerance (T factor): 3

Wind erodibility group (WEG): 3

Wind erodibility index (WEI): 86

Kw factor (surface layer) .20

Land capability, nonirrigated: 3s

Flooding: none

Hydric soil: no
Ponding: none

Hydrologic group: B

Drainage class: somewhat excessively drained Potential for frost action: low

Representative soil profile:	Texture	Permeability	capacity	рН
Ap,A 0 to 14 in	coarse sandy loam	moderately rapid	1.84 to 2.13 in	6.1 to 7.3
Bw 14 to 17 in	coarse sandy loam	moderately rapid	0.30 to 0.39 in	6.1 to 7.3
2Bw.2C 17 to 80 in	gravelly coarse sand	rapid	1.26 to 3.15 in	7.4 to 8.4



Available weter

Sherburne County, Minnesota

# 1224--Hubbard-Verndale complex, 0 to 3 percent slopes

#### Hubbard

Extent: 60 percent of the unit Soil loss tolerance (T factor): 5

Landform(s): stream terraces, outwash plains

Wind erodibility group (WEG): 2

Slope gradient: 0 to 3 percent

Wind erodibility index (WEI): 134

Parent material: outwash

Kw factor (surface layer) .15

Restrictive feature(s): greater than 60 inches

Land capability, nonirrigated: 4s

Flooding: none Hydric soil: no
Ponding: none Hydrologic group: A

Drainage class: excessively drained Potential for frost action: low

Representative soil profile:	Texture	Permeability	capacity	рН
Ap 0 to 11 in	loamy coarse sand	rapid	0.88 to 1.32 in	5.1 to 7.3
Bw 11 to 27 in	loamy sand	rapid	0.48 to 1.13 in	5.1 to 7.3
BC C 27 to 80 in	coarse sand	rapid	1.58 to 3.69 in	5.6 to 7.8

#### Verndale

Extent: 35 percent of the unit Soil loss tolerance (T factor): 3

Landform(s): stream terraces, outwash plains

Wind erodibility group (WEG): 3

Slope gradient: 0 to 2 percent

Wind erodibility index (WEI): 86

Parent material: outwash Kw factor (surface layer) .20

Restrictive feature(s): greater than 60 inches Land capability, nonirrigated: 3s

Flooding: none Hydric soil: no
Ponding: none Hydrologic group: B

Drainage class: somewhat excessively drained Potential for frost action: low

Representative soil prof	file: Texture	Permeability	capacity	рН
Ap 0 to 10 ir	n coarse sandy loam	moderately rapid	1.28 to 1.67 in	5.1 to 7.3
Bt 10 to 16 in	n coarse sandy loam	moderate	0.88 to 1.13 in	5.1 to 7.3
2Bw 16 to 45 in	n coarse sand	rapid	1.72 to 2.30 in	5.1 to 7.3
2C 45 to 80 ir	n sand	rapid	0.70 to 2.10 in	5.1 to 7.3



Available water

Sherburne County, Minnesota

## 1231--Hubbard-Mosford complex, 0 to 3 percent slopes

#### Hubbard

Extent: 60 percent of the unit

Soil loss tolerance (T factor): 5

Landform(s): stream terraces, outwash plainsWind erodibility group (WEG): 2Slope gradient: 0 to 3 percentWind erodibility index (WEI): 134Parent material: outwashKw factor (surface layer) .15Restrictive feature(s): greater than 60 inchesLand capability, nonirrigated: 4s

Flooding: none

Hydric soil: no
Ponding: none

Hydrologic group: A

Drainage class: excessively drained Potential for frost action: low

Representative	soil profile:	Texture	Permeability	Available water capacity	рН
Ap,A	0 to 13 in	loamy sand	rapid	1.04 to 1.56 in	5.1 to 7.3
Bw 1	13 to 19 in	loamy sand	rapid	0.18 to 0.41 in	5.1 to 7.3
BC,C 1	19 to 80 in	sand	rapid	1.83 to 4.27 in	5.6 to 7.8

#### Mosford

Extent: 35 percent of the unit Soil loss tolerance (T factor): 3

Landform(s): stream terraces, outwash plains

Wind erodibility group (WEG): 3

Slope gradient: 0 to 2 percent

Wind erodibility index (WEI): 86

Parent material: outwash Kw factor (surface layer) .20
Restrictive feature(s): greater than 60 inches Land capability, nonirrigated: 3s

Flooding: none

Hydric soil: no

Hydrologic group: B

Drainage class: somewhat excessively drained Potential for frost action: low

Representative soil profile:	Texture	Permeability	capacity	рН
Ap,A 0 to 13 in	sandy loam	moderately rapid	1.69 to 2.34 in	5.1 to 7.3
Bw 13 to 16 in	coarse sandy loam	moderately rapid	0.38 to 0.54 in	5.1 to 7.3
2Bw 16 to 35 in	coarse sand	rapid	0.57 to 2.08 in	5.1 to 7.3
2C 35 to 80 in	sand	rapid	0.90 to 3.14 in	5.1 to 7.8



Available weter

Sherburne County, Minnesota

## 1253B--Stonelake-Sanburn complex, 1 to 6 percent slopes

#### **Stonelake**

Extent: 60 percent of the unit

Landform(s): hills on moraines

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 8

Slope gradient: 1 to 6 percent

Wind erodibility index (WEI): 0

Parent material: outwash

Kw factor (surface layer) .05

Restrictive feature(s): greater than 60 inches

Land capability, nonirrigated: 4s

Flooding: none Hydric soil: no
Ponding: none Hydrologic group: A

Drainage class: excessively drained Potential for frost action: low

Representative soil profil	e: Texture	Permeability	capacity	рН
Ap 0 to 4 in	gravelly loamy sand	rapid	0.24 to 0.59 in	5.1 to 6.5
Bw 4 to 11 in	gravelly coarse sand	very rapid	0.21 to 0.28 in	5.1 to 6.5
Bt 11 to 24 in	very gravelly coarse sand	very rapid	0.39 to 1.04 in	5.1 to 6.5
BC.C 24 to 80 in	gravelly sand	verv rapid	1.12 to 2.80 in	5.1 to 7.8

#### Sanburn

Extent: 30 percent of the unit

Landform(s): hills on moraines

Soil loss tolerance (T factor): 3

Wind erodibility group (WEG): 3

Slope gradient: 1 to 6 percent

Wind erodibility index (WEI): 86

Parent material: outwash

Restrictive feature(s): greater than 60 inches

Land capability, nonirrigated: 3e

Flooding: noneHydric soil: noPonding: noneHydrologic group: B

Drainage class: somewhat excessively drained Potential for frost action: low

Representative	e soil profile:	Texture	Permeability	Available water capacity	рН
Ap	0 to 5 in	sandy loam	moderately rapid	0.61 to 0.77 in	5.1 to 6.5
Bt	5 to 20 in	gravelly sandy loam	moderately rapid	1.05 to 1.80 in	5.1 to 6.5
2BC,2C 2	20 to 80 in	gravelly coarse sand	rapid	1.20 to 2.39 in	5.1 to 6.5



Sherburne County, Minnesota

# 1253C--Stonelake-Sanburn complex, 6 to 15 percent slopes

#### **Stonelake**

Extent: 65 percent of the unit

Landform(s): outwash plains

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 3

Slope gradient: 6 to 15 percent

Wind erodibility index (WEI): 86

Parent material: outwash

Kw factor (surface layer) .02

Restrictive feature(s): greater than 60 inches

Land capability, nonirrigated: 6s

Flooding: none

Hydric soil: no

Ponding: none

Hydrologic group: A

Drainage class: excessively drained Potential for frost action: low

Representative soil prof	ile: Texture	Permeability	capacity	pН
Ap 0 to 4 in	gravelly loamy sand	rapid	0.20 to 0.35 in	5.1 to 6.5
Bw 4 to 11 in	gravelly coarse sand	very rapid	0.07 to 0.64 in	5.1 to 6.5
Bt 11 to 24 in	very gravelly coarse sand	very rapid	0.13 to 0.91 in	5.1 to 6.5
BC. C 24 to 80 in	gravelly sand	very rapid	0.56 to 3.35 in	5.1 to 7.3

#### Sanburn

Extent: 25 percent of the unit

Landform(s): outwash plains

Soil loss tolerance (T factor): 3

Wind erodibility group (WEG): 3

Slope gradient: 6 to 15 percent

Wind erodibility index (WEI): 86

Parent material: outwash

Kw factor (surface layer) .20

Restrictive feature(s): greater than 60 inches

Land capability, nonirrigated: 4s

Flooding: none Hydric soil: no
Ponding: none Hydrologic group: A

Drainage class: somewhat excessively drained Potential for frost action: moderate

Representative	soil profile:	Texture	Permeability	Available water capacity	рН
Ар	0 to 5 in	sandy loam	moderately rapid	0.61 to 0.77 in	5.1 to 6.5
Bt	5 to 20 in	gravelly sandy loam	moderately rapid	1.05 to 1.80 in	5.1 to 6.5
2BC,2C 2	20 to 80 in	gravelly coarse sand	very rapid	1.20 to 2.39 in	5.1 to 6.5



A collecte contact

Sherburne County, Minnesota

# 1253E--Stonelake-Sanburn complex, 15 to 40 percent slopes

#### **Stonelake**

Extent: 65 percent of the unit

Landform(s): hills on moraines

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 2

Slope gradient: 15 to 40 percent

Wind erodibility index (WEI): 134

Parent material: outwash

Kw factor (surface layer) .17

Restrictive feature(s): greater than 60 inches

Land capability, nonirrigated: 7s

Flooding: none

Hydric soil: no

Ponding: none

Hydrologic group: A

Drainage class: excessively drained Potential for frost action: low

Representative	e soil profile:	Texture	Permeability	capacity	рН
Α	0 to 2 in	loamy coarse sand	rapid	0.12 to 0.30 in	5.1 to 6.5
E	2 to 8 in	very gravelly loamy coarse sand	very rapid	0.18 to 0.24 in	5.1 to 6.5
Bt	8 to 16 in	very gravelly coarse sand	very rapid	0.25 to 0.66 in	5.1 to 6.5
C	16 to 80 in	gravelly coarse sand	very rapid	1.28 to 3.19 in	5.1 to 7.8

#### Sanburn

Extent: 25 percent of the unit

Landform(s): hills on moraines

Soil loss tolerance (T factor): 3

Wind erodibility group (WEG): 3

Slope gradient: 15 to 30 percent

Wind erodibility index (WEI): 86

Parent material: outwash

Kw factor (surface layer) .20

Restrictive feature(s): greater than 60 inches

Land capability, nonirrigated: 6e

Flooding: noneHydric soil: noPonding: noneHydrologic group: B

Drainage class: somewhat excessively drained Potential for frost action: low

Representative	soil profile:	Texture	Permeability	Available water capacity	рН
Α	0 to 5 in	sandy loam	moderately rapid	0.61 to 0.77 in	5.1 to 6.5
Bt	5 to 14 in	sandy loam	moderately rapid	0.63 to 1.09 in	5.1 to 6.5
2C 1	14 to 80 in	coarse sand	rapid	1.31 to 2.63 in	5.1 to 6.5



Sherburne County, Minnesota

# 1254--Ricelake fine sandy loam, 0 to 3 percent slopes

#### **Ricelake**

Extent: 90 percent of the unit

Landform(s): outwash plains

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 3

Slope gradient: 0 to 3 percent

Wind erodibility index (WEI): 86

Parent material: glacial outwash over till

Kw factor (surface layer) .20

Restrictive feature(s): greater than 60 inches

Land capability, nonirrigated: 2w

Flooding: none Hydric soil: no
Ponding: none Hydrologic group: B

Drainage class: somewhat poorly drained Potential for frost action: high

Representative soil profile:	Texture	Permeability	Available water capacity	рН
A 0 to 9 in	fine sandy loam	moderately rapid	1.45 to 1.63 in	6.1 to 7.3
E 9 to 27 in	loamy fine sand	rapid	1.06 to 1.95 in	6.1 to 7.3
Bt 27 to 48 in	fine sandy loam	moderately rapid	2.76 to 3.61 in	6.1 to 7.3
2Ca 48 to 80 in	clay loam	moderate	4.46 to 6.06 in	6.1 to 7.8



Sherburne County, Minnesota

# 1255--Elkriver fine sandy loam, 0 to 2 percent slopes, occasionally flooded

## Elkriver, occasionally flooded

Extent: 90 percent of the unit

Landform(s): flood plains

Soil loss tolerance (T factor): 4

Wind erodibility group (WEG): 3

Slope gradient: 0 to 2 percent

Wind erodibility index (WEI): 86

Parent material: alluvium

Kw factor (surface layer) .20

Restrictive feature(s): greater than 60 inches

Land capability, nonirrigated: 2w

Flooding: occasional Hydric soil: no
Ponding: none Hydrologic group: B

Drainage class: somewhat poorly drained Potential for frost action: moderate

Representative soil profile:	Texture	Permeability	Available water capacity	рН
Ap 0 to 10 in	fine sandy loam	moderately rapid	1.57 to 1.97 in	5.1 to 7.3
A1,A3 10 to 26 in	fine sandy loam	moderately rapid	2.42 to 3.23 in	5.1 to 7.3
Bw 26 to 32 in	very fine sandy loam	moderately rapid	0.89 to 1.12 in	5.6 to 7.8
2C 32 to 80 in	sand	rapid	0.96 to 4.80 in	5.6 to 7.8

#### 1256--Cantlin loamy fine sand, 0 to 3 percent slopes

#### Cantlin

Extent: 90 percent of the unit

Landform(s): outwash plains

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 2

Slope gradient: 0 to 3 percent

Wind erodibility index (WEI): 134

Parent material: outwash

Kw factor (surface layer) .17

Restrictive feature(s): greater than 60 inches

Land capability, nonirrigated: 4s

Flooding: none


Hydric soil: no

Ponding: none

Hydrologic group: A

Drainage class: well drained Potential for frost action: low

Representative s	oil profile:	Texture	Permeability	Available water capacity	рН
Ap 0	to 8 in	loamy fine sand	rapid	0.79 to 0.94 in	5.1 to 6.0
Bw 8	to 22 in	loamy fine sand	rapid	0.85 to 1.13 in	5.1 to 6.0
BC,C 22	to 80 in	fine sand	rapid	2.89 to 4.05 in	5.1 to 6.5



Sherburne County, Minnesota

# 1257--Elkriver-Mosford complex, 0 to 6 percent slopes, rarely flooded

#### Elkriver, rarely flooded

Extent: 55 percent of the unit

Landform(s): flood plains

Soil loss tolerance (T factor): 4

Wind erodibility group (WEG): 3

Slope gradient: 0 to 3 percent

Wind erodibility index (WEI): 86

Parent material: alluvium

Kw factor (surface layer) .20

Restrictive feature(s): greater than 60 inches

Land capability, nonirrigated: 2s

Flooding: rare Hydric soil: no Ponding: none Hydrologic group: B

Drainage class: moderately well drained Potential for frost action: moderate

Representative soil profile:	Texture	Permeability	capacity	рН
Ap 0 to 11 in	fine sandy loam	moderately rapid	1.76 to 2.20 in	5.1 to 7.3
AB 11 to 20 in	fine sandy loam	moderately rapid	1.36 to 1.81 in	5.1 to 7.3
Bw 20 to 34 in	fine sandy loam	moderately rapid	2.07 to 2.62 in	5.6 to 7.8
2C 34 to 80 in	fine sand	rapid	0.92 to 4.61 in	5.6 to 7.8

#### Mosford, rarely flooded

Extent: 35 percent of the unit

Landform(s): flood plains

Soil loss tolerance (T factor): 3

Wind erodibility group (WEG): 3

Slope gradient: 0 to 6 percent

Wind erodibility index (WEI): 86

Parent material: outwash

Kw factor (surface layer) .20

Restrictive feature(s): greater than 60 inches

Land capability, nonirrigated: 3e

Flooding: rare


Hydric soil: no

Ponding: none

Hydrologic group: B

Drainage class: somewhat excessively drained Potential for frost action: low

Representative soil profile:	Texture	Permeability	Available water capacity	рН
Ap 0 to 11 in	fine sandy loam	moderately rapid	1.43 to 1.98 in	5.1 to 7.3
Bw 11 to 16 in	fine sandy loam	moderately rapid	0.61 to 0.87 in	5.1 to 7.3
2Bw 16 to 25 in	fine sand	rapid	0.27 to 1.00 in	5.1 to 7.3
2C 25 to 80 in	sand	rapid	1.09 to 3.83 in	5.1 to 7.8



A collecte contact

Sherburne County, Minnesota

# 1258B--Zimmerman fine sand, thick solum, 1 to 6 percent slopes

#### Zimmerman, thick solum

Extent: 95 percent of the unit

Landform(s): hills on outwash plains

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 1

Slope gradient: 1 to 6 percent

Wind erodibility index (WEI): 220

Parent material: outwash

Kw factor (surface layer) .17

Restrictive feature(s): greater than 60 inches

Land capability, nonirrigated: 4s

Flooding: none Hydric soil: no
Ponding: none Hydrologic group: A

Drainage class: excessively drained Potential for frost action: low

Representative soil profile	e: Texture	Permeability	Available water capacity	рН
A 0 to 2 in	fine sand	rapid	0.14 to 0.18 in	5.1 to 6.5
Bw,E 2 to 62 in	fine sand	rapid	3.59 to 5.98 in	5.1 to 6.5
E'&Bt 62 to 80 in	fine sand	rapid	1.09 to 1.81 in	5.1 to 7.3

# 1258C--Zimmerman fine sand, thick solum, 6 to 12 percent slopes

#### Zimmerman, thick solum

Extent: 95 percent of the unit

Landform(s): hills on outwash plains

Soil loss tolerance (T factor): 5

Landform(s): hills on outwash plains

Wind erodibility group (WEG): 1

Slope gradient: 6 to 12 percent

Wind erodibility index (WEI): 220

Parent material: outwash

Kw factor (surface layer) .17


Restrictive feature(s): greater than 60 inches

Land capability, nonirrigated: 6s

Flooding: none Hydric soil: no
Ponding: none Hydrologic group: A

Drainage class: excessively drained Potential for frost action: low

Representativ	ve soil profile:	Texture	Permeability	capacity	pН
Α	0 to 3 in	fine sand	rapid	0.22 to 0.28 in	5.1 to 6.5
Bw,E	3 to 70 in	fine sand	rapid	4.02 to 6.69 in	5.1 to 6.5
E'&Bt	70 to 80 in	fine sand	rapid	0.59 to 0.98 in	5.1 to 7.3



Available water

Sherburne County, Minnesota

# 1258E--Zimmerman fine sand, thick solum, 12 to 35 percent slopes

## Zimmerman, thick solum

Extent: 95 percent of the unit

Landform(s): hills on outwash plains

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 1

Slope gradient: 12 to 35 percent

Wind erodibility index (WEI): 220

Parent material: outwash

Kw factor (surface layer) .17

Restrictive feature(s): greater than 60 inches

Land capability, nonirrigated: 6s

Flooding: none Hydric soil: no
Ponding: none Hydrologic group: A

Drainage class: excessively drained Potential for frost action: low

Representative soil profi	ile: Texture	Permeability	Available water capacity	рН
A 0 to 2 in	fine sand	rapid	0.14 to 0.18 in	5.1 to 6.5
Bw,E 2 to 62 in	fine sand	rapid	3.59 to 5.98 in	5.1 to 6.5
E'&Bt 62 to 80 in	fine sand	rapid	1.09 to 1.81 in	5.1 to 7.3



Sherburne County, Minnesota

## 1260B--Stonelake-Nebish complex, 2 to 6 percent slopes

#### **Stonelake**

Extent: 55 percent of the unit

Landform(s): hills on moraines

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 2

Slope gradient: 2 to 6 percent

Wind erodibility index (WEI): 134

Parent material: outwash

Kw factor (surface layer) .17

Restrictive feature(s): greater than 60 inches

Land capability, nonirrigated: 4s

Flooding: none Hydric soil: no
Ponding: none Hydrologic group: A

Drainage class: excessively drained Potential for frost action: low

Representative	soil profile:	Texture	Permeability	Available water capacity	рН
Ap	0 to 8 in	loamy coarse sand	very rapid	0.79 to 0.94 in	5.1 to 6.5
Bt	8 to 30 in	very gravelly loamy coarse sand	very rapid	0.66 to 1.76 in	5.1 to 6.5
BC.C 3	80 to 80 in	gravelly sand	very rapid	1.00 to 2.50 in	5.1 to 7.4

#### Nebish

Extent: 30 percent of the unit

Landform(s): hills on moraines

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 3

Slope gradient: 2 to 6 percent

Wind erodibility index (WEI): 86

Parent material: till

Kw factor (surface layer) .24

Restrictive feature(s): greater than 60 inches

Land capability, nonirrigated: 2e
Flooding: none

Hydric soil: no

Ponding: none Hydrologic group: B

Drainage class: well drained Potential for frost action: moderate

Representativ	e soil profile:	Texture	Permeability	capacity	рН
Ap	0 to 6 in	fine sandy loam	moderately rapid	0.77 to 1.06 in	5.6 to 7.3
E	6 to 9 in	fine sandy loam	moderately rapid	0.35 to 0.60 in	5.6 to 7.3
Bt	9 to 43 in	clay loam	moderate	5.08 to 6.43 in	5.6 to 7.8
Bk	43 to 80 in	loam	moderate	4.07 to 7.03 in	7.4 to 7.8



Available water

Sherburne County, Minnesota

## 1260C--Stonelake-Nebish complex, 6 to 12 percent slopes

#### **Stonelake**

Extent:55 percent of the unitSoil loss tolerance (T factor):5Landform(s):hills on morainesWind erodibility group (WEG):2Slope gradient:6 to 12 percentWind erodibility index (WEI):134Parent material:outwashKw factor (surface layer).17

Restrictive feature(s): greater than 60 inches Land capability, nonirrigated: 6s

Flooding: none Hydric soil: no
Ponding: none Hydrologic group: A

Drainage class: excessively drained Potential for frost action: low

Representative s	soil profile:	Texture	Permeability	capacity	рН
Ap 0	to 7 in	loamy sand	very rapid	0.71 to 0.85 in	5.1 to 6.5
E 7	to 20 in	loamy coarse sand	very rapid	0.39 to 0.52 in	5.1 to 6.5
Bt 20	to 42 in	very gravelly coarse sand	very rapid	0.66 to 1.76 in	5.1 to 6.5
C 42	to 80 in	very gravelly coarse sand	very rapid	0.76 to 1.89 in	5.1 to 7.8

#### **Nebish**

Extent: 30 percent of the unit

Landform(s): hills on moraines

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 3

Slope gradient: 6 to 12 percent

Wind erodibility index (WEI): 86

Parent material: till

Kw factor (surface layer) .24

Restrictive feature(s): greater than 60 inches

Land capability, nonirrigated: 3e

Flooding: none Hydric soil: no
Ponding: none Hydrologic group: B

Drainage class: well drained Potential for frost action: moderate

Representative soil profile	Texture	Permeability	Available water capacity	рН
Ap 0 to 3 in	fine sandy loam	moderately rapid	0.41 to 0.57 in	5.6 to 7.3
E 3 to 10 in	fine sandy loam	moderately rapid	0.74 to 1.27 in	5.6 to 7.3
Bt 10 to 29 in	clay loam	moderate	2.89 to 3.67 in	5.6 to 7.8
Bk 29 to 80 in	loam	moderate	5.59 to 9.65 in	7.4 to 7.8



Sherburne County, Minnesota

## 1260E--Stonelake-Nebish complex, 12 to 25 percent slopes

#### **Stonelake**

Extent: 60 percent of the unit

Landform(s): hills on moraines

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 8

Slope gradient: 12 to 25 percent

Wind erodibility index (WEI): 0

Parent material: outwash

Kw factor (surface layer) .05

Restrictive feature(s): greater than 60 inches

Land capability, nonirrigated: 7s

Flooding: none

Hydric soil: no

Ponding: none

Hydrologic group: A

Drainage class: excessively drained Potential for frost action: low

Representative soil	profile:	Texture	Permeability	capacity	рН
A 0 to	5 in	gravelly coarse sandy loam	rapid	0.31 to 0.77 in	5.1 to 6.5
Bw 5 to	11 in	very gravelly coarse sand	very rapid	0.18 to 0.24 in	5.1 to 6.5
Bt 11 to	20 in	very gravelly coarse sand	very rapid	0.27 to 0.72 in	5.1 to 6.5
BC.C 20 to	80 in	very gravelly coarse sand	verv rapid	1.20 to 2.99 in	5.1 to 7.8

#### **Nebish**

Extent: 25 percent of the unit

Landform(s): hills on moraines

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 3

Slope gradient: 12 to 25 percent

Wind erodibility index (WEI): 86

Parent material: till

Kw factor (surface layer) .24


Restrictive feature(s): greater than 60 inches

Land capability, nonirrigated: 6e

Flooding: none Hydric soil: no
Ponding: none Hydrologic group: B

Drainage class: well drained Potential for frost action: moderate

Representative	soil profile:	Texture	Permeability	Available water capacity	рН
A 0	) to 5 in	fine sandy loam	moderately rapid	0.67 to 0.92 in	5.6 to 7.3
EB 5	to 9 in	fine sandy loam	moderately rapid	0.43 to 0.75 in	5.6 to 7.3
Bt 9	to 27 in	clay loam	moderate	2.72 to 3.44 in	5.6 to 7.8
Bk 27	7 to 80 in	loam	moderate	5.80 to 10.02 in	7.4 to 7.8



Sherburne County, Minnesota

# 1270B--Milaca fine sandy loam, moderately wet, 3 to 6 percent slopes

# Milaca, moderately wet

Extent: 90 percent of the unit Soil loss tolerance (T factor): 4

Landform(s): drumlins Wind erodibility group (WEG): 3

Slope gradient: 3 to 6 percent Wind erodibility index (WEI): 86

Parent material: till Kw factor (surface layer) .28

Restrictive feature(s): dense material at 40 to 60 inches

Land capability, nonirrigated: 2e

Flooding: none

Hydric soil: no

Ponding: none

Hydrologic group: C

Drainage class: moderately well drained Potential for frost action: moderate

Representative soil profile	Texture	Permeability	capacity	рН
A 0 to 6 in	fine sandy loam	moderately rapid	0.77 to 1.06 in	5.1 to 6.5
E,BE 6 to 19 in	fine sandy loam	moderately rapid	2.34 to 2.86 in	5.1 to 6.5
Bt 19 to 28 in	fine sandy loam	moderate	1.09 to 1.45 in	5.1 to 6.5
BC 28 to 45 in	fine sandy loam	slow	0.00 to 1.35 in	5.6 to 7.3
Cd 45 to 80 in	sandy loam	impermeable	0.00 to 1.40 in	5.6 to 7.3



A collecte contact

Sherburne County, Minnesota

## 1288--Seelyeville-Markey complex, ponded, 0 to 1 percent slopes

## Seelyeville, ponded

Extent: 60 percent of the unit

Soil loss tolerance (T factor): 3

Landform(s): depressions on outwash plains, depressions on

Wind erodibility group (WEG): 8

stream terraces

Slope gradient: 0 to 1 percent

Wind erodibility index (WEI): 0

Parent material: organic material

Kw factor (surface layer) .02

Restrictive feature(s): greater than 60 inches Land capability, nonirrigated: 8w

Flooding: none Hydric soil: yes
Ponding: frequent Hydrologic group: A/D

Drainage class: very poorly drained Potential for frost action: high

Representative soil profile:	Texture	Permeability	Available water capacity	рН
Oa1 0 to 15 in	muck	moderately rapid	5.24 to 6.73 in	
Oa2.Oa3 15 to 80 in	muck	moderately rapid	22.74 to 29.23 in	

#### Markey, ponded

Extent: 30 percent of the unit Soil loss tolerance (T factor): 2

Landform(s): depressions on outwash plains, depressions on Wind erodibility group (WEG): 8 stream terraces

Slope gradient: 0 to 1 percent Wind erodibility index (WEI): 0

Parent material: organic material over outwash

Restrictive feature(s): greater than 60 inches

Kw factor (surface layer) .02


Land capability, nonirrigated: 8w

Flooding: none Hydric soil: yes

Ponding: frequent Hydrologic group: A/D

Drainage class: very poorly drained Potential for frost action: high

Representative soil profile:	Texture	Permeability	capacity	рН
Oa 0 to 27 in	muck	moderately rapid	9.51 to 12.22 in	
A 27 to 32 in	loamy sand	rapid	0.14 to 0.38 in	
Cg 32 to 80 in	sand	rapid	1.44 to 3.84 in	



I Available water

Sherburne County, Minnesota

# 1356--Water, miscellaneous

## Water, miscellaneous

Extent: 100 percent of the unit Soil loss tolerance (T factor):

Landform(s): Wind erodibility group (WEG):
Slope gradient: Wind erodibility index (WEI):
Parent material: Kw factor (surface layer)

Restrictive feature(s): greater than 60 inches

Land capability, nonirrigated:

Flooding: Hydric soil:

Ponding: Hydrologic group:

Drainage class: Potential for frost action:

Representative soil profile:

Texture

Permeability

Available water capacity

pH

Sherburne County, Minnesota

## 1946--Fordum-Winterfield complex, 0 to 2 percent slopes, frequently flooded

# Fordum, frequently flooded

Extent: 65 percent of the unit

Landform(s): flood plains

Soil loss tolerance (T factor): 4

Wind erodibility group (WEG): 8

Slope gradient: 0 to 1 percent

Wind erodibility index (WEI): 0

Parent material: alluvium

Kw factor (surface layer) .20

Restrictive feature(s): greater than 60 inches

Land capability, nonirrigated: 6w

Flooding: frequent Hydric soil: yes
Ponding: none Hydrologic group: D

Drainage class: very poorly drained Potential for frost action: high

Representative	soil profile:	Texture	Permeability	Available water capacity	рН
Α	0 to 7 in	fine sandy loam	moderately rapid	0.78 to 1.28 in	5.1 to 7.3
Cg	7 to 28 in	sandy loam	moderately rapid	2.09 to 4.59 in	5.1 to 7.3
2Cg 28	8 to 80 in	sand	rapid	2.08 to 5.20 in	5.6 to 7.3

#### Winterfield, frequently flooded

Extent: 20 percent of the unit

Landform(s): flood plains

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 8

Slope gradient: 0 to 2 percent

Wind erodibility index (WEI): 0

Parent material: alluvium

Kw factor (surface layer) .17

Restrictive feature(s): greater than 60 inches

Land capability, nonirrigated: 4w

Flooding: frequent

Ponding: none

Hydric soil: no

Hydrologic group: A

Drainage class: somewhat poorly drained Potential for frost action: moderate

Representativ	e soil profile:	Texture	Permeability	Available water capacity	рН
A	0 to 8 in	loamy fine sand	rapid	0.79 to 0.94 in	5.6 to 7.3
C1,C2	8 to 20 in	sand	rapid	0.73 to 1.34 in	5.6 to 7.3
C3,C5	20 to 80 in	sand	rapid	2.39 to 5.98 in	5.6 to 7.3



Sherburne County, Minnesota

#### W--Water

#### Water

Extent: 100 percent of the unit Soil loss tolerance (T factor):

Landform(s): Wind erodibility group (WEG):
Slope gradient: Wind erodibility index (WEI):
Parent material: Kw factor (surface layer)

Restrictive feature(s): greater than 60 inches

Land capability, nonirrigated:

Flooding: Hydric soil:
Ponding: Hydrologic group:

Drainage class: Potential for frost action:

Representative soil profile:

Texture

Permeability

Available water capacity

pH

This report provides a semitabular listing of some soil and site properties and interpretations that are valuable in communicating the concept of a map unit. The report also provides easy access to the commonly used conservation planning information in one place. The major soil components in each map unit are displayed. Minor components may be displayed if they are included in the database and are selected at the time the report is generated.

